


INORGANIC ACID SALTS OF SIBUTRAMINE

Technical Field

The present invention relates to novel inorganic acid salts of sibutramine (N,N-dimethyl-1-[1-(4-chlorophenyl)cyclobutyl]-3-methylbutylamine), represented by Chemical Formula 1, below, and crystalline forms thereof. The present invention is also concerned with pharmaceutical compositions comprising the compounds as effective ingredients, methods of preparing the compounds, and the use of the compounds.

[Chemical Formula 1]



X = HSO₄, Br, H₂PO₄·H₂O

Background Art

Sibutramine (N-[1-[1-(4-chlorophenyl)cyclobutyl]-3-methylbutyl]-N,N-dimethylamine), which is an inhibitor of 5-

hydroxytryptamine and noradrenaline reuptake *in vivo*
(Neuropharmacology, 28, p129-134), is useful in the treatment
of depression, Parkinson's disease, obesity, insulin-
independent diabetes mellitus, epilepsy, and the like. In
5 addition, sibutramine reduces body weight gain by a dual
action to reduce food intake by enhancing satiety and to
increase energy expenditure by stimulating heat generation
(Int. J. Obesity, 19, p145; Brit. J. Pharmacol. 114, p388).

Since sibutramine is difficult to purify due to its
10 low melting point, it is preferable to use a crystalline
material capable of being purified by recrystallization in
order to prepare a pharmaceutical composition comprising
sibutramine. Korean Pat. Publication No. 1990-0000274
discloses that sibutramine is utilized as salts formed with
15 acids providing non-toxic acid addition salts containing
pharmaceutically acceptable anions, for example, in the form
of hydrochloride, malate, acetate, citrate, fumarate,
tartrate, succinate, aspartate or glutmate salt. However,
since sibutramine hydrochloride is difficult to handle
20 pharmaceutically due to its hygroscopic nature, it is
undesirable to use sibutramine hydrochloride for preparing
medicaments. In the preparation of medicaments, a constant
weight of an active compound should be contained in each
dosage form, but an active ingredient absorbing water from
25 the surrounding environment makes it difficult to achieve
such consistency. Korean Pat. Publication No. 94-8913

discloses that when sibutramine hydrochloride is prepared in a monohydrate form, a non-hygroscopic product is obtained, which is suitable for the preparation of capsules, tablets and other pharmaceutical dosage forms.

5 The therapeutic use of sibutramine in depression is described in British Pat. No. 2098602. The therapeutic use of sibutramine in Parkinson's disease is disclosed in International Pat. Publication No. WO88/06444. The therapeutic use of sibutramine in cerebral function disorders
10 is disclosed in U.S. Pat. No. 4,939,175. The use of sibutramine hydrochloride in the treatment of obesity is disclosed in European Pat. No. 397831. Also, International Pat. Publication No. WO95/20949 discloses the use of sibutramine for improving impaired glucose tolerance or
15 glucose tolerance in patients suffering from insulin-independent diabetes mellitus.

 In addition, Brazilian Pat. Publication No. 0105486 discloses a novel salt of sibutramine, sibutramine sulfate, in which two moles of sibutramine are bonded to one mole of
20 sulfuric acid. However, this compound is structurally different from sibutramine hydrogen sulfate (in which one mole of sibutramine is bonded to one mole of sulfuric acid) according to the present invention. In particular, the Brazilian Patent Publication never mentions crystalline
25 forms or physical properties, such as solubility and stability, of the novel salt.

Typically, the preparation of salts having pharmaceutically useful physical properties must satisfy the following physicochemical criteria: (1) good solubility, (2) good stability, (3) good non-hygroscopicity and (4) compressibility into tablet form.

However, Korean Pat. Publication No. 94-8913 states that sibutramine hydrochloride has been known to contain a variable amount of water and thus be hygroscopic, and that non-hygroscopic sibutramine can be obtained by preparing sibutramine hydrochloride in a monohydrate form. Sibutramine hydrochloride monohydrate has been prepared by brining it into contact with a medium consisting of water or a medium containing water.

Thus, sibutramine hydrochloride monohydrate is prepared by a complicated process including adding a predetermined amount of water to a reaction mixture, or including preparing sibutramine hydrochloride anhydrate and suspending the sibutramine hydrochloride anhydrate in a water-containing solvent for a long time with agitation. In addition, since currently available sibutramine hydrochloride monohydrate has relatively low solubility between pH 1.0 and pH 7.4, substitute salts having better solubility need to be developed in order to improve the bioavailability of sibutramine. The term "sibutramine", as used herein, refers to racemic sibutramine, unless otherwise indicated.

Based on this background, the present inventors found

that hydrogen sulfate and bromate salts of sibutramine possess remarkably high solubility in water as well as having non-hygroscopicity and stability, and that sibutramine phosphate hydrate has greatly enhanced solubility even when it exists in a hydrous form, compared to conventional sibutramine hydrochloride hydrate, thereby leading to the present invention.

Disclosure of the Invention

In this regard, intensive and through research into the development of a novel salt of sibutramine, capable of solving the problems encountered in the prior art, conducted by the present inventors, resulted in the finding that inorganic acid salts of sibutramine, particularly hydrogen sulfate, bromate, and phosphate monohydrate, possess good physicochemical properties (solubility, non-hygroscopicity and stability). The present inventors further found that sibutramine anhydrate can be prepared with no additional complicated procedure of adding a predetermined amount of water in order to prepare a hydrous form of sibutramine, and has remarkably high solubility although it is in an anhydrous form, as well as being non-hygroscopic, and that the inorganic acids used are less-toxic acids that have been used in many medicaments, thereby leading to the present invention.

It is therefore an object of the present invention to provide a pharmaceutical composition for treating and preventing pathological states of obesity and related disorders, comprising an inorganic acid salt of sibutramine, which has increased water solubility, is non-hygroscopic, and is stable to heat, as an active ingredient.

It is another object of the present invention to provide the inorganic acid salt of sibutramine, and a method of preparing the same.

It is a further object of the present invention to provide anhydrous crystalline and hydrous crystalline forms of the inorganic acid salt of sibutramine.

It is yet another object of the present invention to provide a pharmaceutical composition comprising the inorganic acid salt of sibutramine as an effective ingredient along with a pharmaceutically acceptable diluent or carrier.

It is still another object of the present invention to provide a method of treating obesity, depression, Parkinson's disease, insulin-independent diabetes mellitus and epilepsy using the inorganic acid salt of sibutramine as an effective ingredient.

Brief Description of the Drawings

The above and other objects, features and other

advantages of the present invention will be more clearly understood from the following detailed description taken in conjunction with the accompanying drawings, in which:

Fig. 1 is an X-ray diffraction spectrum of a first crystalline sibutramine hydrogen sulfate according to Example 1;

Fig. 2 is an X-ray diffraction spectrum of a first crystalline sibutramine hydrogen sulfate according to Example 2;

Fig. 3 is an X-ray diffraction spectrum of a first crystalline sibutramine hydrogen sulfate according to Example 3;

Fig. 4 is an X-ray diffraction spectrum of a first crystalline sibutramine hydrogen sulfate according to Example 4;

Fig. 5 is an X-ray diffraction spectrum of a first crystalline sibutramine hydrogen sulfate according to Example 5;

Fig. 6 is an X-ray diffraction spectrum of a first crystalline sibutramine hydrogen sulfate according to Example 6;

Fig. 7 is an X-ray diffraction spectrum of a second crystalline sibutramine hydrogen sulfate according to Example 7;

Fig. 8 is an X-ray diffraction spectrum of a third crystalline sibutramine hydrogen sulfate according to Example

8;

Fig. 9 is an X-ray diffraction spectrum of crystalline sibutramine bromate according to Example 9; and

Fig. 10 is an X-ray diffraction spectrum of crystalline sibutramine phosphate hydrate according to Example 10.

Best Mode for Carrying Out the Invention

To accomplish the objects of the present invention, the present invention provides inorganic acid salts of sibutramine, preferably crystalline sibutramine hydrogen sulfate and crystalline sibutramine bromate in anhydrous forms, and crystalline sibutramine phosphate monohydrate in a hydrous form.

The present invention also provides a method of preparing an inorganic acid salt of sibutramine, comprising reacting sibutramine with an inorganic acid selected from among sulfuric acid, bromic acid, and phosphoric acid in an inert solvent.

The present invention further provides methods of preparing anhydrous crystalline and hydrous crystalline forms of an inorganic acid salt of sibutramine.

The present invention still further provides a pharmaceutical composition for treating obesity, comprising a therapeutically effective amount of an inorganic acid salt

of sibutramine and a pharmaceutically acceptable diluent or carrier. The present invention provides a method of treating obesity, comprising administering the therapeutically effective amount of the inorganic acid salt
5 of sibutramine.

The present invention still further provides a pharmaceutical composition for treating depression, comprising a therapeutically effective amount of an inorganic acid salt of sibutramine and a pharmaceutically
10 acceptable diluent or carrier. The present invention provides a method of treating depression, comprising administering the therapeutically effective amount of the inorganic acid salt of sibutramine.

The present invention still further provides a pharmaceutical composition for treating Parkinson's
15 disease, comprising a therapeutically effective amount of an inorganic acid salt of sibutramine and a pharmaceutically acceptable diluent or carrier. The present invention provides a method of treating Parkinson's disease,
20 comprising administering the therapeutically effective amount of the inorganic acid salt of sibutramine.

The present invention still further provides a pharmaceutical composition for treating insulin-independent diabetes mellitus, comprising a therapeutically effective
25 amount of an inorganic acid salt of sibutramine and a pharmaceutically acceptable diluent or carrier. The present

invention provides a method of treating insulin-independent diabetes mellitus, comprising administering the therapeutically effective amount of the inorganic acid salt of sibutramine.


5 The present invention still further provides a pharmaceutical composition for treating epilepsy, comprising a therapeutically effective amount of an inorganic acid salt of sibutramine and a pharmaceutically acceptable diluent or carrier. The present invention
10 provides a method of treating epilepsy, comprising administering the therapeutically effective amount of the inorganic acid salt of sibutramine.

The pharmaceutical composition of the present invention is preferably formulated into tablets or capsules.

15 Hereinafter, the present invention will be described in more detail.

The present invention relates to an inorganic acid salt of sibutramine, represented by Chemical Formula 1.

[Chemical Formula 2]



Sibutramine bromate anhydrate according to the present invention displays solubility, non-hygroscopicity, formulability and chemical stability, identical to or better than the commercially available sibutramine hydrochloride monohydrate of Chemical Formula 2. Sibutramine hydrogen sulfate and sibutramine phosphate hydrate exhibit non-hygroscopicity, formulability chemical stability and flowability, identical to or better than sibutramine hydrochloride monohydrate, and in particular exhibit about at least 10 times greater solubility in distilled water and buffer solutions of pH 1.2, pH 4.0, pH 5.3, pH 6.8 and pH 7.4. With respect to non-hygroscopicity, the aforementioned inorganic acid salts of sibutramine display no hygroscopicity and no decrease in water content when they are exposed to relative humidities of 10%, 75% and 90% for a period of seven days or longer. With respect to stability, the inorganic acid salts of sibutramine do not generate impurities and do not change in content even when they are exposed to a high temperature of 60°C for a period of one month or longer. The inorganic acid salts of sibutramine also exhibit good photostability.


The present inventors learned about that the sulfuric, phosphoric and bromic acids, contained in the inorganic acid salts of sibutramine according to the present invention, are typically used in a number of medicaments and are less-toxic acids that have been proven

safe for long-term use, and concluded that the novel inorganic acid salts of sibutramine are suitable for long-term administration, thereby leading to the present invention.

5 The inorganic acid salts of sibutramine according to the present invention may be crystalline or non-crystalline. Crystalline forms of the inorganic acid salts of sibutramine are preferred with respect to physical properties such as non-hygroscopicity and thermodynamical
10 stability.

 The present invention includes a method of preparing the inorganic acid salt of sibutramine. That is, the present invention includes a method of preparing an inorganic acid salt of sibutramine, comprising reacting
15 sibutramine with an inorganic acid in an inert solvent. The reaction using sulfuric acid among inorganic acids used takes place according to the following Reaction 1.

[Reaction 1]



20 Among the inorganic acids used as reactants, sulfuric acid has a reported oral-rat LD₅₀ of 2,140 mg/kg, and has

been used in a number of medicaments, including clopidogrel, cefpirome, amphetamine, salbutamol and gentamycin.

In an embodiment, a first crystalline sibutramine
5 hydrogen sulfate is prepared using acetone, ethyl acetate, ethanol, acetonitrile, methylethyl ketone or methylene chloride as an inert solvent according to the method. This compound is characterized by having an X-ray diffraction pattern in which peaks ($I/I_0 \geq 200$) appear at 2θ values of
10 6.50, 12.18, 12.38, 12.58, 13.06, 14.00, 16.76, 17.04, 18.06, 19.68, 20.32, 20.63, 21.34, 21.82, 22.28, 22.54, 23.32, 24.50, 25.80, 26.42, 28.24, 28.64, 29.28, and 33.34.


In another embodiment, a second crystalline sibutramine hydrogen sulfate is prepared using
15 isopropylether as an inert solvent according to the method. This compound is characterized by having an X-ray diffraction pattern in which peaks ($I/I_0 \geq 100$) appear at 2θ values of 5.73, 6.49, 12.18, 12.51, 13.13, 14.02, 14.79, 16.97, 17.38, 20.62, 21.40, 21.83, 22.31, 22.68, 24.51,
20 24.88, 25.82, 26.45, and 31.60.

In a further embodiment, a third crystalline sibutramine hydrogen sulfate is prepared using methanol and isopropylether as an inert solvent mixture according to the method. This compound is characterized by having an X-ray
25 diffraction pattern in which peaks ($I/I_0 \geq 100$) appear at 2θ values of 6.64, 10.24, 13.03, 15.04, 17.00, 17.53, 17.08,

19.06, 20.52, 22.72, 23.23, 24.23, 25.70, 26.40, and 27.57.

The reaction using bromic acid among inorganic acids used takes place according to the following Reaction 2.

[Reaction 2]




The bromic acid used as a reactant has a reported oral-mouse LD₅₀ of 2,700 mg/kg, and has been used in a number of medicaments, including citalopram, dextromethorphan, fenoterol, galantamine and scopolamine.

10 In detail, the product of Reaction 2, crystalline sibutramine bromate, is characterized by having an X-ray diffraction pattern in which peaks ($I/I_0 \geq 200$) appear at 2θ values of 6.96, 11.48, 13.88, 16.64, 17.14, 18.14, 19.68, 20.92, 21.32, 21.86, 22.16, 22.86, 24.30, 26.16, 26.40,

15 27.42, 28.06, 28.32, 29.52, 31.58, 32.94, 34.54, 37.42, and 37.82.

The reaction using phosphoric acid among inorganic acids used takes place according to the following Reaction 3.

20 [Reaction 3]



The phosphoric acid used as a reactant has a reported oral-rat LD₅₀ of 1,530 mg/kg, and has been used in a number of medicaments, including clindamycin, chloroquine, codeine, disopyramide, metromidazole and oleandomycin. In detail, the product of Reaction 3, crystalline sibutramine phosphate, is characterized by having an X-ray diffraction pattern in which peaks ($I/I_0 \geq 200$) appear at 2θ values of 7.66, 10.68, 11.06, 11.50, 14.46, 15.40, 15.74, 17.22, 17.84, 18.08, 18.98, 19.68, 21.18, 21.50, 21.88, 22.84, 23.18, 23.62, 24.42, 24.72, 25.98, 27.52, 28.38, 28.64, and 29.28.

The inorganic acids used in Reactions 1, 2 and 3 are less-toxic acids that haven been proven safe for long-term use.

The inert solvent available in the preparation method of the present invention includes acetone, ethyl acetate, methanol, ethanol, isopropanol, acetonitrile, isopropyl ether, methylethyl ketone and dichloromethane. Acetone and ethyl acetate are preferred.

In the inert solvent, one equivalent of sibutramine may be reacted with 1 to 2 equivalents, preferably 1.02 to 1.2 equivalents, of concentrated sulfuric acid, at -5 to 40°C, preferably 20 to 30°C, for 0.5 to 5 hours, preferably

2 to 3 hours. Herein, the concentrated sulfuric acid is used after being diluted with the inert solvent.

The preparation method of the present invention may provide an inorganic acid salt of sibutramine in a yield of
5 higher than 90.0% and a high purity of greater than 99.0%.

The present invention provides a pharmaceutical composition for treating or preventing pathological states of obesity and related disorders, comprising a therapeutically effective amount of hydrogen sulfate,
10 bromate or phosphate monohydrate of sibutramine and a pharmaceutically acceptable diluent or carrier, and a method of treating or preventing pathological states of obesity and related disorders by administering this composition.

15 The present invention also provides a pharmaceutical composition for treating depression, comprising a therapeutically effective amount of an inorganic acid salt of sibutramine and a pharmaceutically acceptable diluent or carrier, and a method of treating depression by
20 administering this composition.

The present invention further provides a pharmaceutical composition for treating or preventing Parkinson's disease, comprising a therapeutically effective amount of an inorganic acid salt of sibutramine and a
25 pharmaceutically acceptable diluent or carrier, and a method of treating or preventing Parkinson's disease by

administering this composition.

The present invention still further provides a pharmaceutical composition for treating insulin-independent diabetes mellitus, comprising a therapeutically effective
5 amount of an inorganic acid salt of sibutramine and a pharmaceutically acceptable diluent or carrier, and a method of treating insulin-independent diabetes mellitus by administering this composition.

The present invention still further provides a
10 pharmaceutical composition for treating epilepsy, comprising a therapeutically effective amount of an inorganic acid salt of sibutramine and a pharmaceutically acceptable diluent or carrier, and a method of treating epilepsy by administering this composition.

15 The pharmaceutical composition comprising the inorganic acid salt of sibutramine according to the present invention as an active ingredient may be preferably administered orally, for example in the form of tablets or capsules.

20 Tablets may be prepared by mixing an active ingredient with a carrier, a diluent or an excipient and compressing the mixture into tablets. Examples of suitable carriers, diluents or excipients include 'disintegrators such as starch, sugars and mannitol; fillers and extenders
25 such as calcium phosphate and silicic derivatives; binding agents such as carboxymethyl cellulose and other cellulose

derivatives, gelatin and polyvinyl pyrrolidone; and lubricants such as talc, calcium and magnesium stearate, and solid polyethylene glycol. Also, hard or soft gelatin capsules containing an active ingredient, either with or
5 without an additive such as the carriers, diluents or excipients may be prepared according to an ordinary method.

The pharmaceutical composition preferably contains a crystalline inorganic acid salt of sibutramine, represented by Chemical Formula 1, as an active ingredient in an amount
10 of 1 to 50 parts by weight based on 250 parts by weight of the composition.

For example, the pharmaceutical composition having a total weight of 250 mg according to the present invention may be prepared in such a manner as to contain 10 mg (based
15 on sibutramine content) of the crystalline inorganic acid salt of sibutramine, represented by Chemical Formula 1, 115 mg of microcrystalline cellulose, 115 mg of lactose, 5 mg of silicon dioxide, and 5 mg of magnesium stearate. However, this composition of the pharmaceutical composition
20 is illustrative, and thus, the scope of the present invention is not limited thereto.

A better understanding of the present invention may be obtained through the following examples which are set forth to illustrate, but are not to be construed as the
25 limit of the present invention.

REFERENCE EXAMPLE 1: Preparation of sibutramine hydrochloride monohydrate

Sibutramine hydrochloride anhydrate was prepared according to a method described in Korean Pat. No. 2098602 or Korean Pat. Publication No. 90-00274. Then, according to
5 a method described in British Pat. No. 2184122 or Korean Pat. Publication No. 94-08913, 10 g of the prepared sibutramine hydrochloride anhydrate was dissolved in a boiling mixture of 110 ml acetone and 1.2 ml water, and the
10 resulting solution was hot-filtered and distilled to remove 80 ml of the solvent, thus reducing the volume of the filtrate. The concentrate was filtered to recover a generated solid. The solid was vacuum-dried, thus obtaining 9.2 g (yield: 87%) of the compound of Chemical Formula 2,
15 which had a melting point of 195°C.

EXAMPLES

Hydrogen sulfate, bromate and phosphate monohydrate salts of sibutramine were prepared according to the preparation method of the present invention, and were
20 compared with sibutramine hydrochloride hydrate for physical properties including hygroscopicity, solubility, stability, light stability and crystallizability. In addition, the inorganic acid salts of sibutramine were

formulated into capsules in order to examine their formulability and release patterns.

EXAMPLE 1: Preparation of sibutramine hydrogen sulfate using acetone

5 Sibutramine (17.1 g, 0.06 mol) was dissolved in 150 ml of acetone with agitation. After the solution was adjusted to 25°C, 6.0 g of concentrated sulfuric acid was diluted with 50 ml of acetone and added to the solution. Crystals formed slowly. The resulting mixture was agitated
10 at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 30 ml of acetone, and vacuum-dried at 40°C, thus obtaining 21.0 g (yield: 91%) of a target compound.

TABLE 1

Elemental analysis (C ₁₇ H ₂₈ ClNO ₄ S)	Unit (%)
Measured value	C: 54.35, H: 7.68, N: 3.82, O: 17.00, S: 8.58
Theoretical value	C: 54.03, H: 7.47, N: 3.71, O: 16.93, S: 8.4

15 Melting point (DSC): 212.8°C

¹H-NMR (δ, DMSO-d₆): 8.39 (1H, br, s), 7.54~7.49 (4H, dd), 3.75 (1H, t), 2.83 (3H, d), 2.5 (2H, d), 2.33 (2H, t),

2.13 (3H, d), 1.90 (1H, m), 1.70~1.67 (2H, m), 1.40 (2H, m),
1.00 (6H, t)

**EXAMPLE 2: Preparation of sibutramine hydrogen sulfate
using ethyl acetate**

5 Sibutramine (17.1 g, 0.06 mol) was dissolved in 150
ml of ethyl acetate with agitation. After the solution was
adjusted to 25°C, 6.0 g of concentrated sulfuric acid was
diluted with 50 ml of ethyl acetone and added to the
solution. Crystals formed slowly. The resulting mixture
10 was agitated at 25°C for 2 hrs and further agitated at 4°C
for 1 hr. The generated solid was recovered by filtration
under pressure, washed with 50 ml of ethyl acetate, and
vacuum-dried at 40°C, thus obtaining 21.5 g (yield: 94%) of
a target compound.

15 Melting point: 212°C

**EXAMPLE 3: Preparation of sibutramine hydrogen chloride
using ethanol**

Sibutramine (17.1 g, 0.06 mol) was dissolved in 70 ml
of ethanol with agitation. After the solution was adjusted
20 to 25°C, 6.0 g of concentrated sulfuric acid was diluted

with 10 ml of ethanol and added to the solution to slowly form crystals. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 50 ml of acetone, and vacuum-dried at 40°C, thus obtaining 20.3 g (yield: 89.5%) of a target compound.

Melting point: 211°C

EXAMPLE 4: Preparation of sibutramine hydrogen chloride using acetonitrile

10

Sibutramine (17.1 g, 0.06 mol) was dissolved in 150 ml of acetonitrile with agitation. After the solution was adjusted to 25°C, 6.0 g of concentrated sulfuric acid was diluted with 50 ml of acetonitrile and added to the solution. Crystals formed slowly. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 50 ml of acetonitrile, and vacuum-dried at 40°C, thus obtaining 21.0 g (yield: 92%) of a target compound.

15

20

Melting point: 211°C

EXAMPLE 5: Preparation of sibutramine hydrogen sulfate using methylethyl ketone

Sibutramine (17.1 g, 0.06 mol) was dissolved in 150 ml of methylethyl ketone with agitation. After the solution
5 was adjusted to 25°C, 6.0 g of concentrated sulfuric acid was diluted with 50 ml of methylethyl ketone and added to the solution. Crystals formed slowly. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by
10 filtration under pressure, washed with 50 ml of methylethyl ketone, and vacuum-dried at 40°C, thus obtaining 22.0 g (yield: 97%) of a target compound.

Melting point: 212°C

EXAMPLE 6: Preparation of sibutramine hydrogen sulfate using methylene chloride

Sibutramine (17.1 g, 0.06 mol) was dissolved in 150 ml of methylene chloride with agitation. After the solution was adjusted to 25°C, 6.0 g of concentrated sulfuric acid was diluted with 50 ml of methylene chloride and added to

the solution. Crystals formed slowly. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 50 ml of methylene chloride, and vacuum-dried at 40°C, thus obtaining 20.3 g (yield: 90%) of a target compound.

Melting point: 211°C

EXAMPLE 7: Preparation of sibutramine hydrogen sulfate using isopropyl ether

Sibutramine (17.1 g, 0.06 mol) was dissolved in 150 ml of isopropyl ether with agitation. After the solution was adjusted to 25°C, 6.0 g of concentrated sulfuric acid was diluted with 50 ml of isopropyl ether and added to the solution. Crystals formed immediately after isopropyl ether addition. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 50 ml of isopropyl ether, and vacuum-dried at 40°C, thus obtaining 22.1 g (yield: 97%) of a target compound.

Melting point: 207°C

EXAMPLE 8: Preparation of sibutramine hydrogen sulfate using methanol and isopropyl ether

Sibutramine (17.1 g, 0.06 mol) was dissolved in 50 ml of methanol with agitation. After the solution was adjusted to 25°C, 6.0 g of concentrated sulfuric acid was diluted with 10 ml of methanol and added to the solution. The reaction solution was concentrated from 30 ml to 15 ml under pressure. 150 ml of isopropyl ether was slowly added in droplets to the concentrate for 10 min. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 50 ml of isopropyl ether, and vacuum-dried at 40°C, thus obtaining 20.3 g (yield: 90%) of a target compound.

Melting point: 210°C

EXAMPLE 9: Preparation of sibutramine bromate

Sibutramine (28.0 g, 0.1 mol) was dissolved in 280 ml of ethyl acetate with agitation. After the solution was adjusted to 25°C, 17.2 g of 47% bromic acid was slowly added

in droplets to the solution to form crystals. The resulting mixture was agitated at 25°C for 2 hrs and further agitated at 4°C for 1 hr. The generated solid was recovered by filtration under pressure, washed with 100 ml of ethyl acetate, and vacuum-dried at 40°C, thus obtaining 33.1 g (yield: 92%) of a target compound.

TABLE 2

Elemental analysis (C ₁₇ H ₂₇ BrClN)	Unit (%)
Measured value	C: 56.79, H: 7.77, N: 3.89
Theoretical value	C: 56.60, H: 7.54, N: 3.88

Melting point (DSC): 212.4°C

¹H-NMR (δ, DMSO-d₆): 8.61 (1H, br, s), 7.60~7.48 (4H, dd), 3.80 (1H, t), 2.83 (3H, d), 2.50 (2H, d), 2.32 (2H, t), 2.20 (3H, d), 1.90 (1H, m), 1.77~1.68 (2H, m), 1.39 (2H, m), 1.00 (6H, t)

EXAMPLE 10: Preparation of sibutramine phosphate monohydrate

10 g of sibutramine was dissolved in 100 ml of ethyl acetate with agitation. After the solution was adjusted to 25°C, 4.13 g of 85% phosphoric acid was diluted with 30 ml of ethyl acetate and added in droplets to the solution.

Crystals formed slowly. The resulting mixture was agitated at 25°C for 2 hrs. The generated solid was recovered by filtration under pressure and washed with 30 ml of ethyl acetate. The washed sibutramine phosphate anhydrate was mixed with 120 ml of isopropyl ether, 50 ml of acetone and 1.5 ml of water, agitated at 20-30°C for 18 hrs, filtered, and vacuum-dried, thus obtaining 12.8 g (yield: 90%) of a target compound.

The obtained sibutramine phosphate monohydrate was subjected to elemental analysis and melting point analysis, and the results are as follows.

TABLE 3

Elemental analysis (C ₁₇ H ₃₁ ClNO ₅ P)	Unit (%)
Measured value	C: 51.38, H: 7.69, N: 3.50, O: 19.62
Theoretical value	C: 51.58, H: 7.89, N: 3.54, O: 20.21

Melting point (DSC): 174.1°C

¹H-NMR (δ, DMSO-d₆): 8.14 (1H, br, s), 7.38~7.30 (4H, dd), 3.14 (1H, t), 2.51 (1H, d), 2.45 (1H, d), 2.33 (1H, m), 2.22 (5H, t), 2.18 (3H, t), 1.90 (1H, m), 1.67 (1H, m), 1.56 (1H, m), 1.19 (2H, m), 0.95 (3H, d), 0.89 (3H, d)

EXAMPLE 11: Preparation of capsules containing sibutramine hydrogen sulfate

Ingredients were mixed according to the composition described in Table 4, below, to prepare capsules containing sibutramine hydrogen sulfate.

TABLE 4

Ingredients	Content (per capsule)
Sibutramine hydrogen sulfate	Amount corresponding to 10 mg of sibutramine
Lactose	115 mg
Microcrystalline cellulose	115 mg
Silicon dioxide	5 mg
Magnesium stearate	5 mg

5 The ingredients were mixed and filled into hard capsules using a capsule filling machine (Bosche).

EXAMPLE 12: Preparation of capsules containing sibutramine bromate

10 Ingredients were mixed according to the composition described in Table 5, below, to prepare capsules containing sibutramine bromate.

TABLE 5

Ingredients	Content (per capsule)
Sibutramine bromate	Amount corresponding to 10 mg of sibutramine
Lactose	115 mg
Microcrystalline cellulose	115 mg
Silicon dioxide	5 mg
Magnesium stearate	5 mg

The ingredients were mixed and filled into hard capsules using a capsule filling machine (Bosche).

EXAMPLE 13: Preparation of capsules containing sibutramine phosphate monohydrate

5 Ingredients were mixed according to the composition described in Table 6, below, to prepare capsules containing sibutramine phosphate monohydrate.

TABLE 6

Ingredients	Content (per capsule)
Sibutramine phosphate monohydrate	Amount corresponding to 10 mg of sibutramine
Lactose	115 mg
Microcrystalline cellulose	115 mg
Silicon dioxide	5 mg
Magnesium stearate	5 mg

10 The ingredients were mixed and filled into hard capsules using a capsule filling machine (Bosche).

EXAMPLE 14: Evaluation of hygroscopicity of the inorganic acid salts of sibutramine

15 The inorganic acid salts of sibutramine, prepared in Examples 1, 2 and 3, and sibutramine hydrochloride monohydrate were exposed to humid conditions (75% and 90%

RH) at 25° for a period of three days or one weeks. Then, the water content (K.F. water%) of the samples was measured. The results are given in Table 7, below.

TABLE 7

Storage humidity (relative humidity)		75%		90%	
Storage period	Initial	3 days		1 week	
Sibutramine hydrogen sulfate	0.02%	0.02%	0.01%	0.03%	0.03%
sibutramine bromate	0.09%	0.09%	0.09%	0.08%	0.08%
Sibutramine phosphate hydrate	4.25%	4.24%	4.25%	4.25%	4.26%
Sibutramine HCl hydrate	5.5%	5.49%	5.5%	5.5%	5.49%

5 As shown in Table 7, like sibutramine hydrochloride hydrate, hydrogen sulfate, bromate and phosphate hydrate salts of sibutramine displayed almost no change in water content under humid conditions.

10 **EXAMPLE 15: Evaluation of solubility of the inorganic acid salts of sibutramine**

15 The inorganic acid salts of sibutramine, prepared in Examples 1, 2 and 3, and sibutramine hydrochloride monohydrate were evaluated for solubility in solutions having various pH values. The results are given in Table 8, below. In Table 8, the solubility is expressed as milligrams (mg) of sibutramine dissolved per milliliter (ml) of solution.

TABLE 8

Solvents	Salts of sibutramine				Remarks
	Hydrogen sulfate	Bromate	Phosphate	Hydrochloride	
DW	285	12.78	38.86	26.18	Dissolved at 37°C
pH 1.2	333	12.43	23.92	13.36	
pH 4.0	333	3.64	50	9.58	
pH 5.3	400	8.96	50	6.58	
pH 6.8	370	11.08	24.55	23.14	
pH 7.4	400	12.41	50	9.2	

As shown in Table 8, in distilled water (DW) and buffer solutions at various pH values, hydrogen sulfate and phosphate salts of sibutramine had greatly enhanced solubility compared to sibutramine hydrochloride. These results indicate that these salt forms of sibutramine may have better bioavailability than sibutramine hydrochloride monohydrate.

EXAMPLE 16: Evaluation of stability of the inorganic acid salts of sibutramine

The inorganic acid salts of sibutramine, prepared in Examples 1, 2 and 3, and sibutramine hydrochloride monohydrate were exposed to a stringent 60°C heat treatment. The results are summarized in Table 9, below.

TABLE 9

Storage period	Initial	1 wk	2 wks	4 wks
Hydrogen sulfate	1.000	1.000	0.999	0.999
Bromate	1.000	0.999	1.000	0.999
Phosphate	1.000	1.000	0.999	1.000
Hydrochloride	1.000	1.000	0.999	0.999

HPLC was performed under the following conditions.

Wavelength of UV detection: 225 nm

Column: octadecyl silica gel, C18 (4.6×150 mm, 5 µm)

Mobile phase: ammonium phosphate monohydrate (0.05 M,
5 adjusted to pH 6.0 with phosphoric acid) : acetonitrile =
35 : 65

Flow rate: 1.0 ml/min

As shown in Table 9, like sibutramine hydrochloride
monohydrate, the inorganic acid salts of sibutramine
10 displayed almost no change in content upon stringent 60°C
heat treatment. These results indicate that the hydrogen
sulfate, bromate and phosphate monohydrate salts of
sibutramine, like sibutramine hydrochloride monohydrate,
have good chemical stability at high temperature.

15 **EXAMPLE 17: valuation of light stability of the inorganic
acid salts of sibutramine**

A light stability test was performed as follows. The
inorganic acid salts of sibutramine, prepared in Examples
1, 2 and 3, and sibutramine hydrochloride monohydrate were
20 exposed to fluorescent light at 25°C using a light stability
test chamber suitable for the ICH guideline, for storage

periods of 1, 2 and 4 weeks. The results are given in Table 10, below.

TABLE 10

Storage period	Initial	1 wk	2 wks	4 wks
Hydrogen sulfate	1.000	1.000	1.000	0.999
Bromate	1.000	1.000	0.999	0.999
Phosphate monohydrate	1.000	1.000	0.999	0.999
HCl monohydrate	1.000	1.000	0.999	0.999

As shown in Table 10, when content changes of the inorganic acid salts of sibutramine were analyzed by HPLC in order to determine their light stability, the inorganic acid salts of sibutramine, like sibutramine hydrochloride monohydrate, displayed good light stability.

Industrial Applicability

The hydrogen sulfate, bromate and phosphate monohydrate salts of sibutramine according to the present invention have good physicochemical properties including non-hygroscopicity, solubility, stability, formulability and crystallizability. The hydrogen sulfate and phosphate monohydrate salts of sibutramine exhibit an increased solubility, more than 10 times that of sibutramine hydrochloride hydrate. In addition, due to their non-hygroscopic nature, since the hydrogen sulfate, bromate and phosphate monohydrate salts of sibutramine do not change in

content, they are highly suitable for long-term storage and guarantee consistency suitable for the preparation of pharmaceutical dosage forms.

Moreover, since sulfuric, bromic and phosphoric acids
5 used in the preparation of the novel inorganic acid salts of sibutramine are less-toxic acids that haven been proven to be pharmaceutically safe for long-term use, the inorganic acid salts of sibutramine are useful as novel salts of sibutramine.